

A comparative analysis of standard accretion disks spectra: an application to Ultraluminous X-ray Sources

A. Lorenzin and L. Zampieri

INAF-Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, I-35122 Padova, Italy

Accepted ... Received ...; in original form ...

ABSTRACT

We compare standard models of accretion disks around black holes that include the appropriate zero-torque inner boundary condition and relativistic effects on the emission and propagation of radiation. The comparison is performed adopting the multicolor disk blackbody model (MCD) as reference and looking for the parameter space in which it is in statistical agreement with “more physical” accretion disk models. We find simple ‘recipes’ that can be used for adjusting the estimates of the physical inner radius of the disk, the black hole mass and the accretion rate inferred using the parameters of the MCD fits. We applied these results to four ULXs for which MCD spectral fits of their X-ray soft spectral components have been published and find that, in three cases (NGC 1313 X-1, X-2 and M 81 X-9), the black hole masses inferred for a standard disk around a Schwarzschild black hole are in the interval $\sim 100 - 200 M_{\odot}$. Only if the black hole is maximally rotating are the masses comparable to the much larger values previously derived in the literature.

Key words: accretion, accretion disks — black hole physics — X-rays: binaries

1 INTRODUCTION

Studying the spectra of X-ray binaries is one of the main tools for understanding the physics of these source and determine the properties of the accretion disk and the compact object that they host. From an observational standpoint, Galactic black hole candidates (BHCs) show a number of spectral states, defined in terms of the components in their X-ray spectra and the flux level (e.g. McClintock & Remillard 2006; Remillard & McClintock 2006). In the so called *high state* (Tananbaum et al. 1972; Oda 1977), the spectrum is dominated by a soft, thermal component up to 1-2 keV, with a power-law emerging at higher energies. In the *low state* (Oda et al. 1971; Tananbaum et al. 1972) the soft component is not present and the spectrum is well fitted by a power-law with a cut-off at few tens of keV. Transitions to an *intermediate state* (Méndez & van der Klis 1997; Belloni et al. 1997) or a *very high state* (Miyamoto et al. 1991) are also observed in which both components are present and equally important in terms of the emitted flux. It is worth mentioning that this classification is based also on the timing properties of the sources (e.g. Méndez et al. 1998). It is largely accepted that the soft, thermal component originates from an accretion disk, while the power-law is likely produced by Comptonization of the soft disk photons from a hotter phase of the accretion flow (usually referred to as corona). In these assumptions, in principle the properties of the disk and the physical parameters of the accreting black

hole (BH) can be inferred from the study of the soft component.

Similar spectral states (e.g. Foschini et al. 2002; Feng & Kaaret 2005) and correlated spectral/flux variability (e.g. La Parola 2001; Mucciarelli et al. 2007) are also observed in Ultraluminous X-ray Sources (ULXs), very bright, point-like X-ray sources in the field of nearby galaxies (see, e.g., Fabbiano 2006). In fact, several pieces of evidence indicate that the majority of ULXs are X-ray binaries, chiefly among them the detection of periodic modulations in the X-ray flux (e.g. Kaaret et al. 2006; Fabbiano et al. 2006) and the identification of stellar optical counterparts (e.g. Liu et al. 2002, 2004; Kaaret et al. 2004; Mucciarelli et al. 2005). The spectra of several ULXs show a soft, thermal component, similar to that observed in BHCs, which has been interpreted as emission from an accretion disk, although for some ULXs with spectral curvature above 2-3 keV different spectral models have also been proposed (e.g. Stobbart et al. 2006; Gonalves & Soria 2006; Mizuno et al. 2007).

To model the soft component in BHCs and ULXs the so called multicolor disk blackbody model (MCD; Mitsuda et al. 1984) has been often adopted because it is easy to use and is efficiently implemented in the X-ray spectral fitting package XSPEC. The fitting parameters of the MCD model (the disk inner temperature and the normalization of the spectrum) depend on the BH mass and accretion rate, and can in principle be used to estimate the physical parameters of the accretion flow and of the BH. However, it is well

known that this model represents an approximation of the standard, Shakura & Syunyaev (1973) disk and differs from it for the expression of the temperature profile, in which the effects of the viscous torque at the inner boundary are neglected. Also, the locally emitted spectrum is assumed to be a blackbody, neglecting the vertical temperature structure and atmospheric radiative transfer effects. Although the effects of a zero-torque inner boundary in a pseudo-Newtonian potential (Paczynsky & Wiita 1980) were implemented in the *diskpn* model (Gierlinski et al. 1999), temperature profile correction factors for the MCD model were also derived (e.g. Kubota et al. 1998; Makishima et al. 2000). At the same time, limited deviations from LTE caused by radiative transfer effects were accounted for using an hardening factor $f = T_{col}/T_{eff}$, where T_{col} and T_{eff} are the color and effective temperatures (e.g. Shimura & Takahara 1995; Zampieri et al. 2001). Even adopting all these correction factors, deviations of the MCD with respect to a standard disk spectrum may be significant if the physical state of the disk changes or the coupling between the disk and the corona becomes important, which may occur especially at high accretion rates (Merloni et al. 2000).

Despite all these uncertainties, the parameters of the MCD fit have been used for estimating the mass of the BH and the accretion rate in X-ray binary systems. This has recently been done also for ULXs for which, at present, X-ray spectroscopy provides one of the few available methods to obtain zero-th order estimates of the BH mass. Indeed, if the soft component observed in ULX spectra is fitted with a MCD model (e.g. Miller et al. 2003, 2004a), its characteristic temperature is typically ~ 200 eV, much lower than that observed in Galactic BHCs. Both the low temperature and high normalization constant of the MCD model have been used to estimate the BH masses, typically obtaining values largely in excess of $100 M_{\odot}$. Modelling the soft component in these terms is clearly possible if ULXs are in an accretion stage similar to that of BHCs in the soft (or nearly soft) state, so that a disk spectral component is present. Along with the very high (isotropic) luminosity of these sources, these “spectroscopic estimates” have led to the suggestion that they may contain intermediate mass black holes (IMBHs).

In this paper we revisit and compare standard models of accretion disks around black holes that include the appropriate zero-torque inner boundary condition and/or relativistic effects on the emission and propagation of radiation. They are derived in the usual assumptions that: (1) the accretion disk is optically thick, geometrically thin and in a quasi-steady state; (2) the locally emitted spectrum from a small annulus of the disk is a blackbody ($I_{\nu} = B_{\nu}$), with a color correction factor f taking into account for atmospheric transfer effects. The comparison is performed adopting the MCD as reference model and looking for the parameter space in which it turns out to be in agreement with “more physical” accretion disk models. The fits are performed keeping either the BH mass or the accretion rate fixed and allowing the inner radius of the MCD to vary. This gives us simple ‘recipes’ that can be used for adjusting the estimates of the physical inner radius of the disk, the black hole mass and the accretion rate inferred using the parameters of the MCD fits.

The paper is organized as follows. Section 2 contains a short summary of the basic structure equations of the accre-

tion disk models considered in this work; Section 3 presents the results of the comparison among the different spectral models; Section 4 is devoted to an application of the results of this comparison to some representative ULX spectra, while Section 5 summarizes our conclusions.

2 ACCRETION DISK SPECTRA

In this section we summarize the basic equations of the accretion disk models that we have considered in our work. We start with recalling the main properties of the standard disk model (Shakura & Syunyaev 1973) and the multicolor disk blackbody model (MCD, Mitsuda et al. 1984), that are usually adopted in fitting the spectra of X-ray binaries. Then, we shortly review the structure equations for a relativistic accretion disk orbiting around a Schwarzschild and a Kerr black hole (Bardeen et al. 1972; Novikov & Thorne 1973; Page & Thorne 1974). The emergent spectrum is calculated taking into account the relativistic effects on the emission and propagation of light (gravitational redshift, light bending, Doppler shift) using the formalism of the transfer function introduced by Cunningham (1975). More sophisticated treatments of photon propagation in curved space-times, both numerical (Laor 1991; Karas et al. 1992; Bromley et al. 1997; Fanton et al. 1997; Wilms 1998; Rauch & Blandford 1994; Muller & Camenzind 2004; Beckwith & Done 2004) and also analytical (Cadez & Gomboc 1996; Gomboc 2001; Cadez et al. 2003; Calvani 2004) are available. Such techniques adopt ray tracing of the photon geodesics from the disk plane to the observer and are particularly suited when dealing with the calculation of spectral line profiles. However, we are essentially interested in comparing the continuum spectral shape of the MCD disk with that of a standard and relativistic disk and, to this end, a treatment of the relativistic geometrical optics à la Cunningham (1975) appears acceptable.

2.1 Standard accretion disk and MCD

Let’s consider the *Standard accretion disk* model (Shakura & Syunyaev 1973) in the usual approximations (Newtonian, optically thick, geometrically thin and in a quasi-steady state). The expression of the temperature profile is obtained setting the energy rate of viscous dissipation equal to the total radiation flux (e.g. Frank, King & Raine 2002):

$$T(r) = T_* \left(\frac{r}{R_{in}} \right)^{-3/4} \left[1 - \left(\frac{R_{in}}{r} \right)^{1/2} \right]^{1/4}, \quad (1)$$

where

$$T_* = \left(\frac{3GM\dot{M}}{8\pi\sigma R_{in}^3} \right)^{1/4}, \quad (2)$$

M is the black hole mass, $\dot{M}$ is the accretion rate and R_{in} is the inner disk radius. The emitted spectrum is calculated assuming that the disk emits locally as a blackbody, so that:

$$L_{\nu} = 4\pi r_0^2 F_{\nu} = 8\pi^2 \cos i \int_{R_{in}}^{R_{out}} I_{\nu}(T) r dr, \quad (3)$$

where i is the disk's inclination angle with respect to the direction of the line of sight, r_0 is the distance to the observer, R_{out} is the outer disk radius and $I_\nu(T)$ is the specific intensity. We assume that the gas is in LTE and hence $I_\nu(T) = B_\nu(T)$, where $B_\nu(T)$ is the Planck function.

The MCD model is an approximation of the standard disk and differs from it only for the expression of the temperature profile, in which the effects of the viscous torque at the inner boundary are neglected. The temperature profile is then given by (Mitsuda et al. 1984):

$$T(r) = T_{in} \left(\frac{r}{R_{in,BB}} \right)^{-3/4}, \quad (4)$$

where

$$T_{in} = \left(\frac{3GM\dot{M}}{8\pi\sigma R_{in,BB}^3} \right)^{1/4}, \quad (5)$$

and $R_{in,BB}$ is the inner radius of the MCD. The emitted spectrum can again be calculated from equation (3), with the temperature profile given by equation (4).

In the following, we will use the MCD model as reference for fitting the standard and relativistic disk spectra, because this model is frequently adopted for the spectral fits of X-ray binaries. The fitting parameters of the MCD model, the disk inner temperature T_{in} and the spectral normalization factor $K_{BB} = \cos i (R_{in,BB}/r_0)^2$, depend on M and $\dot{M}$. Assuming that the inner radius $R_{in,BB} = bR_g$ ($R_g = GM/c^2$ is the gravitational radius), it is possible to write M and $\dot{M}$ as:

$$\frac{M}{M_\odot} = \frac{67.5}{b} \left(\frac{r_0}{1 \text{ Mpc}} \right) \left(\frac{K_{BB}}{\cos i} \right)^{1/2} \quad (6)$$

$$\frac{\dot{M}}{\dot{M}_{Edd}} = 0.1b^2 \left(\frac{r_0}{1 \text{ Mpc}} \right) \left(\frac{K_{BB}}{\cos i} \right)^{1/2} \left(\frac{T_{in}}{1 \text{ keV}} \right)^4, \quad (7)$$

where $K_{BB} = [(R_{in,BB}/1 \text{ km})^2 (r_0/10 \text{ kpc})^{-2} \cdot \cos i]$.

In these assumptions, if the distance r_0 is known and the inclination angle i is fixed, the MCD parameters can be used for estimating M and/or $\dot{M}$. These expressions depend on the value of b , the inner radius in units of R_g , and T_{in} . It is well known that these estimates of the BH mass and the disk accretion rate are uncertain, because the physical state of the disk may vary and, in addition, radiative transfer effects induced by scattering may cause departures of I_ν from B_ν . Several non-LTE accretion disk models, in which radiative transfer effects at the surface of the disk are computed, have been presented in the literature, for both hot disks around stellar mass BHs (e.g. Merloni et al. 2000; Davis et al. 2005) and cool disks around intermediate mass BHs (e.g. Hui et al. 2005). These effects are very important in predicting the local spectra emitted at a given disk annulus and can be accounted for using an hardening factor $f = T_{col}/T_{eff}$, where T_{col} and T_{eff} are the color and effective temperature. It is easy to show that $K_{BB} \propto 1/f^4$ and, as a consequence, $M \propto f^2$ and $\dot{M}/\dot{M}_{Edd} \propto f^2$ (Kubota et al. 1998, 2001). So, as a first approximation, we account for radiative transfer effects simply inserting f^2 into equations (6) and (7). Computations of f for standard and relativistic disks give values in the interval 1.4-2 (see Shimura & Takahara 1995; Zampieri et al. 2001; Hui et al. 2005; Davis et al. 2005).

2.2 Relativistic accretion disk

The structure equations and the emitted spectrum of the relativistic disk are obtained in the same assumptions stated above: (1) the accretion disk is optically thick, geometrically thin and in a quasi-steady state; (2) the local emitted spectrum from a small annulus of the disk is a blackbody ($I_\nu = B_\nu$).

In the following, we will refer to the radius of the innermost stable circular orbit around a black hole with r_{ms} . Its expression is given by:

$$r_{ms} = M \left\{ 3 + Z_2 - [(3 - Z_1)(3 + Z_1 + 2Z_2)]^{1/2} \right\}, \quad (8)$$

where Z_1 and Z_2 are functions of M and a (Novikov & Thorne 1973), and a is the specific angular momentum of the black hole.

2.2.1 Temperature profile of the relativistic disk

A detailed derivation of the structure of a relativistic accretion disk around a black hole is presented in Novikov & Thorne (1973) and Page & Thorne (1974) to which we refer for all the details. Here we present only the relevant equations for the temperature profile and the locally emitted flux.

Following Novikov & Thorne (1973), the emitted radiation flux can be expressed as a function of the integrated $\phi-r$ component of the stress-energy tensor (evaluated in the comoving frame of the observer in geodetic circular motion) as:

$$F = \frac{3GM\dot{M}}{8\pi r^3} \frac{Q}{BC^{1/2}}, \quad (9)$$

where B , C and Q depend on the radial coordinate r and the specific angular momentum of the black hole a (Novikov & Thorne 1973). The function Q depends also on the angular momentum per unit mass of a generic circular orbit and of the innermost stable circular orbit (Stoeger 1976). Assuming that the radiative transport of energy is dominant over the turbulent one, in LTE we can write

$$F = \sigma T^4. \quad (10)$$

Comparing equations (9) and (10) it is possible to obtain the surface temperature profile for a relativistic accretion disk:

$$T(r) = T_* \left(\frac{r}{R_{in}} \right)^{-3/4} B^{-1/4} C^{-1/8} Q^{1/4}, \quad (11)$$

where T_* is given by equation (2).

2.2.2 Relativistic effects on the propagation of radiation

Following Bardeen et al. (1972) and Cunningham (1975) the spectrum of a relativistic accretion disk is calculated by means of a transfer function f , that accounts for all relativistic effects on the emission and propagation of radiation (Doppler shift, light bending and gravitational redshift). The expression for f is (Cunningham 1975):

$$f(g^*, r_e, \vartheta_0) dg^* dr_e = \frac{g}{\pi r_e} \left(g^* - g^{*2} \right)^{1/2} r_0^2 \cos \vartheta_0 d\Omega_0, \quad (12)$$

where r_e is the radius of a generic emitting ring of the disk, ϑ_0 is the polar angle between the line of sight of the distant

observer and the disk's polar axis, $d\Omega_0$ is the solid angle that a specific geodesics family subtends to the observer at a given energy and for a given emission radius, $g = E_0/E_e$ is the ratio of the observed photon energy and the emitted one, g^* , g_{max} and g_{min} are functions of g (for their expression see Cunningham 1975).

The radiation flux seen by a distant observer is:

$$F_0 = \int I_0 \cos \vartheta_0 d\Omega_0, \quad (13)$$

where I_0 is the observed specific intensity. If I_e is the emitted specific intensity, from the expression of the relativistic invariant I/E^3 and equation (13) we obtain the observed luminosity per unit energy

$$L_0 = 4\pi r_0^2 F_0 = 4\pi \int g^3 I_e r_0^2 \cos \vartheta_0 d\Omega_0. \quad (14)$$

Using equation (12) it is possible to rewrite $r_0^2 \cos \vartheta_0 d\Omega_0$ in terms of $f(g^*, r_e, \vartheta_0) dg^* dr_e$. Inserting the resulting expression into equation (14), we obtain:

$$L_0 = \int 2\pi f g^2 I_e (g^* - g^{*2})^{-1/2} dg^* d(\pi r_e^2). \quad (15)$$

Tabulated values of the transfer function for a Schwarzschild and a maximally rotating ($a = 0.9981$) Kerr BH were reported by Cunningham (1975). In the following we will consider these two extreme cases as reference.

3 PARAMETER ESTIMATES FROM SPECTRAL FITS

We implemented the structure equations and spectra of the accretion disk models outlined above and produced fits of the standard and relativistic disks against the MCD model. This model-model comparison has been performed by means of a standard χ^2 minimization procedure, that provides also an assessment of the statistical significance of the fit. A fixed reference error of $\sim 10\%$ was assigned to the spectra. As mentioned above, we use the MCD model as reference for fitting all the other disk spectra, because this model is frequently adopted for the spectral fits of X-ray binaries within XSPEC. The values of the best-fitting MCD parameters are then compared with those of the other models.

From now on we refer to the different accretion disk models with the following expressions: *Disk-stand* for the standard accretion disk (Shakura & Syunyaev 1973), *Disk-BB* for the MCD (Mitsuda et al. 1984), *Disk-rel* or *Disk-kerr* for a relativistic disk around a Schwarzschild or a maximally rotating ($a = 9981$) Kerr black hole (Novikov & Thorne 1973), respectively. We will also use different indices to denote the mass M and the accretion rate $\dot{M}$ of the different models (M_{BB} , $\dot{M}_{BB}$ for the *Disk-BB*; M_{stand} , $\dot{M}_{stand}$ for the *Disk-stand*; M_{rel} , $\dot{M}_{rel}$ for the *Disk-rel*; M_{kerr} , $\dot{M}_{kerr}$ for the *Disk-kerr*). The accretion rate is expressed in units of the Eddington accretion rate $\dot{M}_{Edd} = L_{edd}/c^2$, where L_{edd} is the Eddington luminosity. For an accretion efficiency $\eta = 0.1 - 0.4$, the accretion luminosity reaches L_{edd} for $\dot{M} = \dot{M}_{Edd}/\eta = 2.5 - 10\dot{M}_{Edd}$.




Figure 1. Spectrum (*top panel*) and temperature profile (*bottom panel*) of the *Disk-stand* model (*solid-green*) for $M_{stand} = 100M_\odot$, $\dot{M}_{stand} = 0.1\dot{M}_{Edd}$, $R_{in} = 6R_g$, and the corresponding best-fitting *Disk-BB* model (*dashed-red*; $M_{BB} = 94M_\odot$, $\dot{M}_{BB} = 0.09\dot{M}_{Edd}$, $R_{in,BB} = 13.9R_g$). These plots refer to the fitting results reported in Table 1, second row ($\cos i = 1$).

3.1 *Disk-stand* model

Fits of the standard disk with a MCD were performed in the literature in order to obtain correction factors for the inner disk radius, the black hole mass and the accretion rate (e.g. Kubota et al. 1998; Makishima et al. 2000). We performed similar fits of the *Disk-stand* with the *Disk-BB* but keeping either the BH mass or the accretion rate fixed. Acceptable fits with different best-fit values of $R_{in,BB}$ were obtained (see Table 1). For the fit with fixed $\dot{M}$, we obtain $R_{in,BB} = 13R_g$, while for the fit with fixed M , $R_{in,BB} = 14.9R_g$. In the first case, the value of $\dot{M}_{BB}$ returned by the fit is $\approx 20\%$ smaller than $\dot{M}_{stand}$ while, in the second case, the values of M_{BB} derived from the fit is $\approx 12\%$ smaller than the corresponding value of M_{stand} . Clearly, there are also fits with fixed $R_{in,BB}$ in the interval $13 \leq R_{in,BB} \leq 14.9$, that return intermediate values of M_{BB} and $\dot{M}_{BB}$ (see again Table 1). In Figure 1 we show the results of one of these fits.

These results are in agreement with those of Kubota et al. (1998), who showed that the inner radius of the best-fitting MCD model does not correspond to the true inner radius of the standard disk, but that the two radii are related by the expression (see also Zimmerman et al. 2005)




Figure 2. Same as Figure 1 for the *Disk-rel* model (solid-orange; $M_{rel} = 100M_{\odot}$, $\dot{M}_{rel} = 0.1\dot{M}_{Edd}$, $R_{in,rel} = 6R_g$). The parameters of the best-fitting *Disk-BB* model (dashed-red) are $M_{BB} = 87M_{\odot}$, $\dot{M}_{BB} = 0.07\dot{M}_{Edd}$, $R_{in,BB} = 21.6R_g$ (see Table 2, fifth row). For both models $\cos i = 1$.

$$R_{in,BB} = 2.6R_{in}. \quad (16)$$

In fact, in order for the spectral peaks of the *Disk-BB* and *Disk-stand* be superimposed (see Figure 1), the maximum temperature of the standard disk ($0.488T_*$; see equations [1] and [2]) must be equal to the maximum temperature of the *Disk-BB* (T_{in} ; see equation [5]). From this, equation (16) easily follows. Therefore, assuming that the *Disk-stand* terminates at the ISCO (innermost stable circular orbit), $R_{in} = 6R_g$ and the inner radius of the best-fitting *Disk-BB* must be located at $\simeq 15.6R_g$, in substantial agreement with the results of the model-model fits.

We note that all the results presented here hold for any inclination angle, as the *Disk-BB* and *Disk-stand* models have the same dependence on i .

3.2 *Disk-rel* model

Similar fits of the *Disk-rel* model with the *Disk-BB* were also performed. For sufficiently small inclination angles, also in this case the emitted spectrum can be well reproduced by the MCD. The fits were performed keeping either M or $\dot{M}$ fixed. Results are shown in Table 2. The two fits return different values of the inner radius, $R_{in,BB} = 19.2R_g$ for




Figure 3. Same as Figure 1 for the *Disk-kerr* model (solid-blue; $M_{kerr} = 100M_{\odot}$, $\dot{M}_{kerr} = 0.1\dot{M}_{Edd}$). The parameters of the best-fitting *Disk-BB* model (dashed-red) are $M_{BB} = 88M_{\odot}$, $\dot{M}_{BB} = 0.095\dot{M}_{Edd}$, $R_{in,BB} = 8.9R_g$ (see Table 3, second row). For both models $\cos i = 1$.

fixed M and $R_{in,BB} = 25R_g$ for fixed $\dot{M}$. In the first case the value of $\dot{M}_{BB}$ returned by the fit is $\approx 40\%$ smaller than $\dot{M}_{rel}$ whereas, in the second case, the value of M_{BB} derived from the fit is $\approx 25\%$ smaller than the corresponding value of M_{rel} . From Table 2 we can see that good fits can be obtained also fixing $R_{in,BB}$ in the interval $19.2\text{--}25R_g$ and leaving M_{BB} and $\dot{M}_{BB}$ free to vary. In Figure 2 we plot the results of one of these fits ($R_{in,BB} \approx 21.6R_g$).

If we increase the inclination angle, the relativistic effects (especially the gravitational focusing) tend to increase the contribution of radiation coming from the inner parts of the disk, making the spectrum of the *Disk-rel* intrinsically harder than that of the *Disk-BB*. In fact, for high inclination angles the *Disk-BB* is only in rough agreement (within a $\sim 10\%$ fractional error) with the *Disk-rel* spectrum, and no satisfactory fit is obtained ($\chi^2_{red} \geq 1.3$). Similar results for $\cos i < 1$ were obtained by Ebisawa et al. (1991). For illustrative purposes alone, in Table 2 we report also the values of the fits for $\cos i = 0.5$.

3.3 *Disk-kerr* model

The last comparison concerns the *Disk-kerr*/*Disk-BB* models. If the inclination angle is close to $i = 0$ (face-on disk),

Table 1. Values of the characteristic parameters of the *Disk-stand* model and those derived by the spectral fits with the *Disk-BB* model for $\cos i = 1$. The inner radius of the *Disk-stand* model is $6R_g$. In the first row M_{BB} is set equal to M_{stand} ; in the second row $R_{in,BB}$ is fixed at an intermediate value between 13 and $14.9 R_g$; in the last row $\dot{M}_{BB}$ is set equal to $\dot{M}_{stand}$. The results of the spectral fit at fixed radius are shown in Figure 1.

M_{stand} ($M_\odot$)	$\dot{M}_{stand}$ ($M_\odot$)	M_{BB} ($\dot{M}_{Edd}$)	$\dot{M}_{BB}$ ($\dot{M}_{Edd}$)	$R_{in,BB}$ (R_g)	χ^2_{red}
100.0	0.1	100.0	0.08 ± 0.01	13.0 ± 1.5	0.1
100.0	0.1	94.0 ± 10.0	0.09 ± 0.03	13.9	0.2
100.0	0.1	88.0 ± 32.0	0.1	14.9 ± 1.3	0.2

Table 2. Values of the characteristic parameters of the *Disk-rel* model and those derived by the spectral fits with the *Disk-BB* model for $\cos i = 1$ and $\cos i = 0.5$. The inner radius of the *Disk-rel* is $R_{in,rel} = 6R_g$. In the first row M_{BB} is set equal to M_{rel} ; in the second row $R_{in,BB}$ is fixed at an intermediate value between 19.2 and $25 R_g$; in the last row $\dot{M}_{BB}$ is set equal to $\dot{M}_{rel}$ (for each value of $\cos i$). The results of the spectral fit at fixed radius are shown in Figure 2.

	M_{rel} ($M_\odot$)	$\dot{M}_{rel}$ ($M_\odot$)	M_{BB} ($\dot{M}_{Edd}$)	$\dot{M}_{BB}$ ($\dot{M}_{Edd}$)	$R_{in,BB}$ (R_g)	χ^2_{red}
$\cos i = 1$	50.0	0.1	50.0	0.06 ± 0.03	19.2 ± 11.6	0.2
	50.0	0.1	43.5 ± 6.7	0.07 ± 0.02	21.6	0.1
	50.0	0.1	37.3 ± 10.2	0.1	25.0 ± 7.1	0.1
$\cos i = 1$	100.0	0.1	100.0	0.06 ± 0.03	19.2 ± 3.5	0.2
	100.0	0.1	87.0 ± 31.0	0.07 ± 0.02	21.6	0.1
	100.0	0.1	75.0 ± 34.0	0.1	25.0 ± 6.7	0.1
$\cos i = 0.5$	100.0	0.1	100.0	0.03 ± 0.05	11.2 ± 3.4	1.3
	100.0	0.1	50.0 ± 32.0	0.1	21.5 ± 12.3	1.8

the spectral fits of the *Disk-kerr* with the *Disk-BB* at constant M or $\dot{M}$ are statistically acceptable and return different values of the inner radius, $R_{in,BB} = 7.9R_g$ and $R_{in,BB} = 9.2R_g$, respectively (see Table 3). The values of $\dot{M}_{BB}$ and M_{BB} are larger than the corresponding values of the *Disk-rel* model (as high as $\sim 20\%$ for $\dot{M}$ and $\sim 25\%$ for M). Also in this case, there are other fits with fixed $R_{in,BB}$ in the range $7.9\text{--}9.2R_g$ for which the *Disk-BB* and *Disk-kerr* spectra are in good agreement (see again Table 3). In Figure 3 we show the results of one of these fits obtained for $R_{in,BB} \approx 8.9R_g$.

We have also tried fits varying the inclination angle i . For a maximally rotating Kerr black hole, the relativistic effects on the emitted spectrum are significantly stronger than for a Schwarzschild black hole. The spectrum of the *Disk-kerr* model becomes intrinsically different and much harder than the *Disk-BB* one as i increases, so that it is no longer possible to obtain a satisfactory fit ($\chi^2 > 2$). For illustrative purposes alone, in Table 3 we report the values of the fits for $\cos i = 0.5$.

3.4 Comparison with fitting results from XSPEC

In order to test our approach, we simulated disk spectra for both a Schwarzschild BH and a maximally rotating Kerr

BH using the *kerrBB* model in XSPEC (v. 12.3.0). This model implements the relativistic structure equations and the relativistic effects on the emission and propagation of radiation (Li et al. 2005). We fitted the *kerrBB* spectra with a *diskBB* model (the XSPEC implementation of the MCD) and used the parameters of the fit to estimate the mass and accretion rate using equations (6) and (7).

Spectra were simulated using the response matrices of the EPIC-pn camera onboard *XMM-Newton*. For the sake of comparison, in the *kerrBB* model we adopted a zero torque inner boundary condition and switched off the effects of self-irradiation and limb-darkening. The disk was assumed to be face-on ($i = 0$) and the color correction factor f was set equal to 1. A fixed distance of 5 Mpc was assumed. For the mass of the BH and the accretion rate we chose $100M_\odot$ and $0.485\dot{M}_{Edd}$, respectively, while the BH angular momentum a was set equal to 0 (Schwarzschild) or 0.9981 (Kerr).

For a *kerrBB* spectrum with $a = 0$, the best fitting MCD parameters are: $T_{in} = 0.122 \pm 0.002$ keV and $K = 39 \pm 6$ (2σ errors). From these parameters, assuming $b = 19.2 \pm 3.5$ (see Table 2), we obtain $M_{BB} = (110 \pm 29)M_\odot$ (eq. [6]), in good agreement with the assumed value of the BH mass. Similarly, assuming $b = 25 \pm 6.7$, it is $\dot{M}_{BB} = (0.4 \pm 0.28)\dot{M}_{Edd}$ (eq. [7]), which is also in agreement with the assumed value for the accretion rate. For a *kerrBB* spec-

Table 3. Values of the characteristic parameters of the *Disk-kerr* model and those derived by the spectral fits with the *Disk-BB* model for $\cos i = 1$ and $\cos i = 0.5$. The inner radius of the *Disk-kerr* model has been calculated using equation (8) (for $a = 0.9981$, $R_{in,kerr} \cong 1.36R_g$). In the first row M_{BB} is set equal to $\dot{M}_{Kerr}$; in the second row $R_{in,BB}$ is fixed at an intermediate value between 7.9 and 9.2 R_g ; in the last row $\dot{M}_{BB}$ is set equal to $\dot{M}_{Kerr}$ (for each value of $\cos i$). The results of the spectral fit at fixed radius are shown in Figure 3.

	M_{Kerr} ($M_\odot$)	$\dot{M}_{Kerr}$ ($M_\odot$)	M_{BB} ($\dot{M}_{Edd}$)	$\dot{M}_{BB}$ ($\dot{M}_{Edd}$)	$R_{in,BB}$ (R_g)	χ^2_{red}
$\cos i = 1$	100.0	0.1	100.0	0.075 ± 0.015	7.9 ± 0.9	0.2
	100.0	0.1	88.0 ± 18.0	0.095 ± 0.02	8.9	0.2
	100.0	0.1	85.0 ± 20.0	0.1	9.2 ± 1.3	0.2
$\cos i = 0.5$	100.0	0.1	100.0	0.04 ± 0.015	2.4 ± 1.7	2.3
	100.0	0.1	60.0 ± 5.0	0.1	4.0 ± 0.4	2.2

Table 4. Values of the BH mass M and accretion rate $\dot{M}$ for NGC 1313 X-1 and NGC 1313 X-2, obtained using the parameters of the MCD that best-fits their soft components ($K_{BB} = 28^{+5}_{-5}$, $T_{in} = 0.22$ keV for NGC 1313 X-1 ; $K_{BB} = 6.66^{+16}_{-3.5}$, $T_{in} = 0.25$ keV for NGC 1313 X-2) and different values of b that correspond to the different accretion disk models (see Tables 2, 3 and section 4; Miller et al. 2003 and Miller et al. 2004a adopted $b = 9.5$). The assumed distance of NGC 1313 is 3.7 Mpc (Tully 1988) and the color correction factor is $f = 1.7$ (Shimura & Takahara 1995; Zampieri et al. 2001; Hui et al. 2005).

	NGC 1313 X-1			NGC 1313 X-2	
$R_{in,BB}$ (R_g)	M ($M_\odot$)	$\dot{M}$ ($\dot{M}_{Edd}$)	M ($M_\odot$)	$\dot{M}$ ($\dot{M}_{Edd}$)	
6	630^{+50}_{-60}	$0.5^{+0.01}_{-0.1}$	310^{+260}_{-100}	$0.4^{+0.3}_{-0.1}$	
7.9	480^{+40}_{-40}	—	240^{+190}_{-80}	—	
9.2	—	$1.1^{+0.1}_{-0.1}$	—	$0.9^{+0.8}_{-0.2}$	
9.5	400^{+40}_{-40}	$1.2^{+0.1}_{-0.1}$	200^{+160}_{-70}	$1.0^{+0.8}_{-0.3}$	
13	290^{+30}_{-30}	—	145^{+120}_{-50}	—	
14.9	—	$3.0^{+0.2}_{-0.4}$	—	$2.4^{+2.0}_{-0.8}$	
19.2	200^{+20}_{-20}	—	100^{+80}_{-30}	—	
25	—	$8.3^{+0.6}_{-0.8}$	—	$6.7^{+5.7}_{-2.0}$	

trum with $a = 0.9981$, the best fitting MCD parameters are: $T_{in} = 0.248 \pm 0.003$ keV and $K = 6 \pm 0.4$ (2σ errors). From equations (6) and (7) and the values of b reported in Table 3, we obtain $M_{BB} = (105 \pm 15)M_\odot$ and $\dot{M}_{BB} = (0.4 \pm 0.14)\dot{M}_{Edd}$, again in satisfactory agreement with the parameters assumed in simulating the spectrum.

4 ESTIMATES OF M AND $\dot{M}$ FROM THE X-RAY SOFT COMPONENT OF ULXS

A certain number of ULXs show two component X-ray spectra that can be adequately modelled with a soft thermal component plus a power-law. If the soft component is fitted with a MCD, the fitting parameters (in particular the normalization K_{BB}) return very large values for the BH mass, often in excess of several hundreds $M_\odot$ (e.g. Miller et al. 2003, 2004a). These masses are calculated from equation (6) assuming specific values for the inner radius b (in units of

R_g) and the color correction factor f . Miller et al. (2003) and Miller et al. (2004a) adopted $b = 9.5$ and $f = 1.7$.

In the previous Section, we performed fits of the standard and relativistic disk with a MCD keeping the BH mass fixed and determined the (range of) values of b for which the spectra are in statistical agreement. These values are reported in Tables 1, 2 and 3 and differ from what assumed by Miller et al. (2003) and Miller et al. (2004a). For the standard disk, there appears to be more agreement with what reported by Kubota et al. (1998) and Makishima et al. (2000). We are then in the position to revise previous estimates of the BH mass (and accretion rate) of ULXs based on MCD spectral fits of the soft component, adopting the appropriate value of b for both a standard and a relativistic disk.

For illustrative purposes, we use the results of the MCD spectral fits of NGC 1313 X-1, NGC 1313 X-2, M 81 X-9 and NGC 4559 X-7 obtained by Miller et al. (2003, 2004a),

Table 5. Values of the BH mass M and accretion rate $\dot{M}$ for M81 X-9 and NGC 4559 X-7, obtained using the parameters of the MCD that best-fits their soft components ($K_{BB} = 20^{+20}_{-10}$, $T_{in} = 0.26$ keV for M81 X-9 (obs. 1); $K_{BB} = 60^{+70}_{-40}$, $T_{in} = 0.21$ keV for M81 X-9 (obs. 2); $K_{BB} = 158^{+340}_{-107}$, $T_{in} = 0.148$ keV for NGC 4559 X-7) and different values of b that correspond to the different accretion disk models (see Tables 2, 3 and section 4; Miller et al. 2003 and Miller et al. 2004a adopted $b = 9.5$). The assumed distance is 9.69 Mpc for NGC 4559 X-7 (Sanders et al. 2003) and 3.4 Mpc for M81 X-9 (Georgiev et al. 1991; Hill 1993). The color correction factor is $f = 1.7$ (Shimura & Takahara 1995; Zampieri et al. 2001; Hui et al. 2005).

M 81 X-9 (obs. 1)			M 81 X-9 (obs. 2)		NGC 4559 X-7	
$R_{in,BB}$ (R_g)	M ($M_\odot$)	$\dot{M}$ ($\dot{M}_{Edd}$)	M ($M_\odot$)	$\dot{M}$ ($\dot{M}_{Edd}$)	M ($M_\odot$)	$\dot{M}$ ($\dot{M}_{Edd}$)
6	490^{+210}_{-140}	$0.7^{+0.3}_{-0.2}$	860^{+400}_{-370}	$0.6^{+0.3}_{-0.2}$	4000^{+3000}_{-1750}	$0.6^{+0.5}_{-0.3}$
7.9	380^{+150}_{-110}	—	650^{+300}_{-280}	—	3000^{+2300}_{-1300}	—
9.2	—	$1.7^{+0.7}_{-0.5}$	—	$1.3^{+0.5}_{-0.6}$	—	$1.4^{+1.1}_{-0.6}$
9.5	310^{+130}_{-90}	$1.8^{+0.8}_{-0.5}$	540^{+260}_{-230}	$1.4^{+0.6}_{-0.6}$	2500^{+1950}_{-1100}	$1.5^{+1.2}_{-0.6}$
13	230^{+90}_{-70}	—	400^{+180}_{-170}	—	1800^{+1400}_{-750}	—
14.9	—	$4.5^{+1.8}_{-1.3}$	—	$3.3^{+1.5}_{-1.4}$	—	$3.8^{+2.9}_{-1.6}$
19.2	160^{+60}_{-50}	—	270^{+120}_{-120}	—	1200^{+1000}_{-500}	—
25	—	$12.5^{+5.2}_{-5.6}$	—	$9.3^{+4.3}_{-3.9}$	—	$10.1^{+8.2}_{-4.6}$

Zampieri et al. (2004) and Cropper et al. (2004). From equations (6) and (7) of Section 2.1, we calculate the values of M and $\dot{M}$ for different values of b and assuming $f = 1.7$, and report them in Tables 4 and 5. The values that correspond to $b = 9.5$ are those reported by Miller et al. (2003) and Miller et al. (2004a), whereas the values for $b = 7.9, 13, 19.2$ correspond to those expected for a standard disk around a maximally rotating Kerr, newtonian, or Schwarzschild BH, respectively. As can be seen from the Tables, the values of the mass are systematically lower (up to a factor 3) than those of a MCD with $R_{in,BB}$ at the ISCO ($b = 6$). This is clearly a consequence of the larger fitting radius required by the newtonian and relativistic disks. Comparing the masses of the different “physical” models, it is possible to see that the largest value is obtained for a disk around a maximally rotating Kerr BH (*Disk-kerr*; see Tables 4 and 5). This is consistent with the fact that the (Boyer-Lindquist) radial coordinate of the ISCO for a maximally rotating, Kerr BH is $\simeq 1/6$ that of a Schwarzschild BH of the same mass (cmp. Hui & Krolik 2008). Indeed, for a fixed value of $R_{in,BB}$ (in cm) as returned by the MCD fit of an observed spectrum, a lower value of b necessarily implies a larger mass as $b = R_{in,BB}/R_g \propto 1/M$. We need to fill the same physical radius $R_{in,BB}$ with a bigger BH in order to reduce the value of b . Ultimately, the smaller value of b returned by the fit of the *Disk-kerr* model is a consequence of the fact that the Kerr spectrum is significantly harder than those of the other “physical” disk models computed with the same value of the parameters (see Figure 4).

It is also interesting to note that, apart from NGC 4559 X-7, the BH masses for a Schwarzschild disk are no longer significantly in excess of $\sim 100M_\odot$, with a distribution clustering in the interval 100-200 $M_\odot$ (see again Tables 4 and 5). This is true also for M 81 X-9, for which the second observation returns somewhat larger values of the mass. However, as

the limits inferred from the two observations of this source must be consistent within the errors, the mass of the BH must be in the interval $150 - 220M_\odot$ (smaller/larger values would not be consistent with the second/first observation; see the last row in Table 5). All the mass estimates obtained for the *Disk-rel* model are then significantly lower than those reported by Miller et al. (2003) and Miller et al. (2004a). The only exception is NGC 4559 X-7 for which, however, there are hints of timing features in the power density spectrum and it has been proposed that it may contain a BH of several hundreds solar masses (e.g. Cropper et al. 2004). The inferred BH masses can reach the values estimated by Miller et al. (2004a) only for a disk around a maximally rotating, Kerr BH.

5 CONCLUSIONS

The spectrum of a standard disk can be well reproduced by a MCD model. However, as already noted by Kubota et al. (1998, 2001, 2005), if one tries to exploit the MCD model to derive the values of the accretion rate and black hole mass, with the usual assumption that, in the soft state, the disk terminates at the ISCO ($R_{in,BB} = 6R_g$), the inferred values of M and $\dot{M}$ are largely overestimated. On the other hand, the spectrum of the standard disk can be fitted by a MCD model with the same values of M or $\dot{M}$ if we take $R_{in,BB} = bR_g$ with $b \simeq 13$ or $b \simeq 15$, respectively. This is in substantial agreement with the approximate value ($b = 15.6$) that can be derived using analytic arguments.

Also in the case of a relativistic disk around a Schwarzschild black hole, for sufficiently small inclination angles the emitted spectrum can be well reproduced by a MCD model (see also Ebisawa et al. 1991 and Kubota et al. 2005). For small angles, in fact, one can try to guess the disk and black hole parameters using the best-fitting MCD




Figure 4. Emitted spectra (*top panel*) and temperature profiles (*bottom panel*) of the *Disk-BB*, *Disk-stand*, *Disk-rel* and *Disk-kerr* models, obtained for the same characteristic parameters: $M = 100M_{\odot}$, $\dot{M} = 0.1\dot{M}_{Edd}$, $R_{in,BB} = R_{in} = R_{in,rel} = 6R_g$, while $R_{in,kerr}$ has been obtained using equation (8). For all models $\cos i = 1$.

model but, as for the standard disk, a suitable assumption must be made on the inner disk radius. We found that there exist precise values of $R_{in,BB}$ ($\simeq 19.2R_g$ and $\simeq 25R_g$) for which the values of M and $\dot{M}$ inferred from the MCD are the same as those provided by the relativistic disk. For large angles, owing to genuine relativistic effects, no satisfactory MCD spectral fit of the relativistic disk spectrum can be obtained (e.g. Ebisawa et al. 1991).

The spectrum of an accretion disk around a fast rotating Kerr black hole, is in general more strongly affected by relativistic effects compared with those of a disk around a Schwarzschild black hole. As a consequence, it can be well reproduced by a MCD model, only in the case in which the disk is essentially face-on. In this assumption, there exist values of $R_{in,BB}$ for which the MCD and Kerr disks return similar values of M and $\dot{M}$ ($R_{in,BB} \simeq 7.9R_g$ and $R_{in,BB} \simeq 9.2R_g$, respectively). Also in this case, for large inclination angles, the spectrum of the *Disk-kerr* model becomes intrinsically different and much harder than the *Disk-BB* one, so that it is no longer possible to obtain a satisfactory fit with the MCD model.

The model-model comparison approach adopted in this Paper was prompted by the idea to determine the parameter

space in which the MCD is in agreement with “more physical” standard accretion disk models and provide “recipes” for adjusting the estimates of the disk inner radius, the black hole mass and the accretion rate. In fact, despite the uncertainties related to radiative transfer and to the physical state of the inner disk, the parameters of the MCD fit are often used for inferring the mass and accretion rate in X-ray binaries, in particular in ULXs. The results of the comparison that we performed can be used to revise estimates of M and $\dot{M}$ obtained in this way. This has also been tested by applying our “recipe procedure” to the results from a *diskBB* fit to simulated *kerrBB* spectra performed directly in XSPEC.

We considered the case of a few ULXs (NGC 1313 X-1, NGC 1313 X-2, M81 X-9 and NGC 4559 X-7) for which MCD spectral fits of their X-ray soft spectral components have been published and/or values of the BH mass estimated (Cropper et al. 2004; Miller et al. 2003, 2004a; Zampieri et al. 2004). From the parameters of the fit with the *Disk-BB* (T_{in} and K_{BB}) we found that, assuming that the inner disk boundary of the MCD is at or close to $6R_g$, the values of M and $\dot{M}$ can be severely overestimated. Adopting the appropriate value of the inner radius b (in units of R_g) for the *Disk-rel* model, we obtain that the BH masses are in the range $\approx 100 - 200M_{\odot}$ for NGC 1313 X-1, NGC 1313 X-2, M81 X-9, and $700 - 2000M_{\odot}$ for NGC 4559 X-7 (see Tables 4 and 5). The relativistic disk models have BH masses systematically lower (up to a factor 3) than those of a MCD with $R_{in,BB}$ at the ISCO.

As already mentioned, sophisticated relativistic accretion disk models (e.g. Hanawa 1989; Li et al. 2005; Hui et al. 2005) that implement the relativistic structure equations and the relativistic effects on the emission and propagation of radiation, are available in XSPEC and can clearly be used to perform accurate fits of observed spectra, in particular for large inclination angles when the relativistic effects produce significant differences. An analysis of this type has recently been performed by Winter et al. (2007) and Hui & Krolik (2008). It is interesting to note that the BH masses for a sample of disk-dominated ULXs obtained by Hui & Krolik (2008) are below $\sim 100M_{\odot}$. Even if their best fits require large values of the BH spin, BHs of several hundreds solar masses are not needed, in substantial agreement with our findings.

Clearly, it is possible that at least some ULXs for which curvature above $\sim 2-3$ keV is present in the X-ray spectrum are in a different accretion regime, in which the disk is in a diverse physical state (accreting at or above the Eddington rate; see e.g. Roberts 2007; Soria 2007). In this case, the assumptions of a standard accretion disk break down and different spectral models should be adopted (e.g. Stobart et al. 2006; Mizuno et al. 2007). However, in the assumption that they are in a standard accretion regime (below Eddington), we found that, in three cases (NGC 1313 X-1, X-2 and M 81 X-9), the black hole masses inferred for a standard disk around a Schwarzschild black hole are in the interval $\sim 100 - 200M_{\odot}$. Only if the black hole is maximally rotating are the masses comparable to the much larger values previously derived in the literature.

6 ACKNOWLEDGEMENTS

We are indebted with an anonymous referee for his/her valuable comments. We thank Andrea Martocchia for his careful reading of the manuscript. LZ acknowledges financial support from INAF through grant PRIN-2007-26.

REFERENCES

- Bardeen, J. M., Press, W. H., Teukolsky, S. A., 1972, *ApJ*, 178, 347
- Beckwith, K., Done, C., 2004, *PThPS*, 155, 259
- Belloni, T., van der Klis, M., Lewin, W. H. G., van Paradijs, J., Dotani, T., Mitsuda, K., & Miyamoto, S. 1997, *A&A*, 322, 857
- Bromley, B. C., Chen, K., Miller, W. A., 1997, *ApJ*, 475, 57
- Cadez, A., Gomboc, A., 1996, *A&A*, 119, 293
- Cadez, A., Brajnik, M., Gomboc, A., Calvani, M., Fanton, C., 2003, *A&A*, 403, 29
- Calvani, M., 2005, *MNRAS*, 363, 177
- Cropper, M., Soria, R., Mushotzky, R. F., Wu, K., Markwardt, C. B., Pakull, M., 2004, *MNRAS*, 349, 39-51
- Cunningham, C. T., 1975, *ApJ*, 202, 788
- Davis, W. S., Blaes, M. O., Turner, J. N. 2005, *ApJ*, 621, 372
- Ebisawa, K., Kazuhisa M., 1991, *ApJ*, 367, 213-220
- Fabbiano, G., 2006, *ARA&A*, 44, 323
- Fabbiano, G. et al. 2006, *ApJ*, 650, 879
- Fanton, C., Calvani, M., de Felice, F., Cadez, A., 1997, *PASJ*, 49, 159
- Feng, H., Kaaret, P., 2005, *ApJ*, 633, 1052
- Foschini, L., Di Cocco, G., Ho, L. C., Bassani, L., Cappi, M., Dadina, M., Gianotti, F., Malaguti, G., Panessa, F., Piconcelli, E., and 2 coauthors, 2002, *A&A*, 392, 817
- Frank, J., King, A., & Raine, D. J. 2002, *Accretion Power in Astrophysics*, pp. 398. ISBN 0521620538. Cambridge, UK: Cambridge University Press, February 2002.
- Georgiev, T. B., Bilkina, B. I., Tikhonov, N. A., & Karachentsev, I. 1991, *A&AS*, 89, 529
- Gierlinski, M., Zdziarski, A. A., Poutanen, J., Coppi, P. S., Ebisawa, K., Johnson, W. N., 1999, *MNRAS*, 309, 496
- Gomboc, A., 2001, *PhDT*, 69
- Gonçalves, A. C., Soria, R., 2006, *MNRAS*, 371, 673
- Hanawa, T., 1989, *ApJ*, 341, 948
- Hui, Y. & Krolik, J. H. 2008, *ApJ*, in press (arXiv:0803.3607)
- Hui, Y., Krolik, J. H., & Hubeny, I. 2005, *ApJ*, 625, 913
- Hill, J. K., et al. 1993, *ApJ*, 402, L45
- Kaaret, P., Ward M. J., Zezas, A., 2004, *MNRAS*, 351, L83
- Kaaret, P., Feng, H., 2006, (arXiv:0707.2055)
- Karas, V., Vokrouhlický, D., Polnarev, A. G., 1992, *MNRAS*, 259, 569
- Kubota, A., Tanaka, Y., Makishima, K., Ueda, Y., Dotani, T., Inoue, H., Yamaoka, K., 1998, *PASJ*, 50, 667
- Kubota, A., Makishima, K., 2001, *ApJ*, 560, L147-L150
- Kubota, A., Ken, E., Makishima, K., Kazuhiro N., 2005, *ApJ*, 631, 1062-1071
- Laor, A., 1991, *ApJ*, 376, 90
- Li, Li-Xin, Zimmerman, E. R., Narayan, R., McClintock, J. E., 2005, *ApJS*, 157, 335
- La Parola V., 2001, *ApJ*, 556, 47
- Liu, Ji-F., Bregman, J. N., Seitzer, P., 2004, *ApJ*, 580, L31
- Liu, Ji-F., Bregman, Joel N., Seitzer, P., 2004, *ApJ*, 602, 249
- Makishima, K. et al. 2000, *ApJ*, 535, 632
- McClintock, J. E., & Remillard, R. A. 2006, in *Compact Stellar X-ray Sources*, ed. W. H. G. Lewin & M. van der Klis (Cambridge: Cambridge Univ. Press), 157
- Méndez, M., & van der Klis, M. 1997, *ApJ*, 479, 926
- Méndez, M., Belloni, T., & van der Klis, M. 1998, *ApJ*, 499, L187
- Merloni, A., Fabian, A. C., Ross, R. R., 2000, *MNRAS*, 313, 193
- Miller, J. M., Fabbiano, G., Miller, M. C., Fabian, A. C., 2003, *ApJ*, 585, 37
- Miller, J. M., Fabian, A. C., Miller, M. C., 2004, *ApJ*, 607, 931-938
- Miller, J. M., Fabian, A. C., Miller, M. C., 2004, *ApJ*, 614, L117-L120
- Mitsuda, K., Inoue, H., Koyama, K., Makishima, K., Matsuoka, M., Ogawara, Y., Suzuki, K., Tanaka, Y., Shibazaki, N., Hirano, T., 1984, *PASJ*, 36, 741
- Miyamoto, S., Kimura, K., Kitamoto, S., Dotani, T., & Ebisawa, K. 1991, *ApJ*, 383, 784
- Mizuno, T. et al. *PASJ*, 59, 257
- Mucciarelli, P., Zampieri, L., Falomo, R., Turolla, R., Treves, A., 2005, *ApJ*, 633, 101
- Mucciarelli, P., Zampieri, L., Treves, A., Turolla, R., Falomo, R., 2007, *ApJ*, 658, 999
- Muller, A., Camenzind, M., 2004, *A&A*, 413, 861
- Novikov, I. D., Thorne, K. S., 1973, in: "Black holes (Les astres occlus)", p. 343, 450
- Oda, M. 1977, *Space Sci. Rev.*, 20, 757
- Oda, M., Gorenstein, P., Gursky, H., Kellogg, E., Schreier, E., Tananbaum, H., & Giacconi, R. 1971, *ApJ*, 166, L1
- Paczynsky, B., Wiita, P. J., 1980, *A&A*, 88, 23
- Page, D. N., Thorne, K. S., 1974, *ApJ*, 191, 499
- Rauch, K. P., Blandford, R. D., 1994, *ApJ*, 421, 46
- Remillard, R. A., & McClintock, J. E. 2006, *ARA&A*, 44, 49
- Roberts, P. T., 2007, *Ap&SS*, in press (arXiv:0706.2562)
- Sanders D. B., Mazzarella J. M., Kim D.-C., Surace J. A., Soifer B. T., 2003, *AJ*, 126, 1607
- Soria, R., 2007, *Ap&SS*, in press (arXiv:0707.2049)
- Shakura, N. I., Syunyaev, R. A., 1973, *A&A*, 24, 337
- Shimura, T., Takahara, F., 1995, *ApJ*, 445, 780
- Stobart, A.-M., Roberts, T. P., Wilms, J., 2006, *MNRAS*, 368, 397
- Stoeger, W. R. 1976, *A&A*, 53, 267
- Tananbaum, H., Gursky, H., Kellogg, E., Giacconi, R., & Jones, C. 1972, *ApJ*, 177, L5
- Tully, R. B., 1988, *Nearby Galaxies Catalog* (Cambridge: Cambridge Univ. Press)
- Wilms, J., Speith, R., Reynolds, C. S., 1998, *LNP*, 514, 69
- Winter, L. M., Mushotzky, R. F., Reynolds, C. S., 2007, *ApJ*, 655, 163
- Zampieri, L., Turolla, R., E. Szuszkiewicz, 2001, *MNRAS*, 325, L266Z
- Zampieri, L., Mucciarelli, P., Falomo, R., Kaaret, P., Di Stefano, R., Turolla, R., Chierigato, M., Treves, A., 2004, *ApJ*, 603, 523
- Zimmerman, E. R., Narayan, R., McClintock, J. E., J. M.

Miller, 2005, ApJ, 618, 832